

# Hypnotherapy, Relaxation, and Music Therapy in Pediatric Cancer Pain Management: A Clinical Trial Comparison



Fayegh Yousefi<sup>1</sup>, Borhan Moradveisi<sup>2</sup>, Daem Roshani<sup>3</sup>, Majid Mansouri<sup>4</sup> and Karo Servatyari<sup>5,\*</sup>




<sup>1</sup>Department of Clinical Psychology, Medical Faculty, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>2</sup>Department of Pediatrics, School of Medicine Cancer and Immunology Research Center, Besat Hospital, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>3</sup>Department of Epidemiology and Biostatistics Social Determinants of Health Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>4</sup>Department of Pediatrics, School of Medicine, Besat Hospital, Kurdistan University of Medical Sciences.

<sup>5</sup>Student Research Committee, Kurdistan University of Medical Science, Sanandaj, Iran



**Fig. (1).** CONSORT DIAGRAM.

Allocated to intervention (n=42) including: Music Therapy (n=13), Hypnotherapy (n=16) and Relaxation Therapy (n=13).

◆ Received allocated intervention (n=42).

◆ Did not receive allocated intervention (give reasons) (n=16).